

Recent Electroweak Results from the Tevatron

Junjie Zhu¹

State University of New York at Stony Brook, Stony Brook, NY, 11794

Abstract. I present the recent electroweak measurements related to single W , Z boson and diboson productions from the CDF and DØ experiments at the Fermilab Tevatron collider.

Keywords: Electroweak measurements, W and Z bosons, diboson processes

PACS: 12.15.-y, 13.38.Be, 14.70.Fm, 13.85.Qk, 14.60.Cd, 13.40.Em, 14.70.Hp, 13.85.Qk

INTRODUCTION

W and Z bosons are mainly produced via quark-antiquark annihilations at the Fermilab Tevatron collider. Precision measurements with these gauge bosons provide us with high precision tests of the Standard Model (SM) as well as indirect search for possible new physics beyond the SM.

W BOSON PROPERTIES

Precision measurement of W boson mass (M_W). In the SM, M_W can be calculated using the electromagnetic coupling constant α , the Fermi constant G_F and the weak mixing angle $\sin^2 \theta_W$. It also receives quantum radiative corrections that depend on the top quark mass and the SM Higgs boson mass. A precise measurement of M_W thus can be used to make constraints on the Higgs mass. To make the equal contribution to the Higgs mass uncertainty, we need to have $\Delta M_W \approx 0.006 \times \Delta M_{top}$. With the current world average value of $\Delta M_{top} = 1.3$ GeV and $\Delta M_W = 0.025$ GeV, ΔM_W is the limiting factor for the Higgs mass constraint.

At the Tevatron, M_W is extracted from a template fit to the transverse mass (M_T), lepton transverse momentum (p_T^ℓ) and missing transverse energy ($\cancel{E}_T$) distributions in $W \rightarrow \ell \nu (\ell = e, \mu)$ events. The template distributions are generated using a parameterized Monte Carlo (MC) simulation program. This program uses the state-of-the-art W and Z MC event generator and simulates the complex detector acceptance and response effects. The parameters used in the simulation are mainly determined using the $Z \rightarrow \ell \ell$ events. The uncertainties on these detector smearing parameters and theoretical calculations (including QCD and electroweak corrections and also the parton distribution functions (PDFs)) are propagated to the uncertainty on the measurement of M_W .

Using 1 fb⁻¹ of Run II data [1], the DØ collaboration measured $M_W = 80.401 \pm 0.021(\text{stat}) \pm 0.038(\text{syst}) = 80.401 \pm 0.043$ GeV, the most precise measurement from

¹ For the CDF and DØ Collaborations




FIGURE 1. The M_T and p_T^e distributions for data and MC simulation with backgrounds added (top), and the difference between data and MC divided by the data uncertainty (χ) for each bin (bottom).

one single experiment to date. Figure 1 shows the data and MC comparison for M_T and p_T^e distributions.

Many of the systematic uncertainties in these measurements are limited by the size of the control samples (mainly $Z \rightarrow \ell\ell$ events) used to understand the detector and physics effects. These uncertainties can be improved with an analysis of the larger datasets in hand. CDF has begun analyzing a data sample with $\mathcal{L} = 2.3 \text{ fb}^{-1}$ [2] and found the statistical uncertainty is 16 (15) MeV for muon (electron) channel and the uncertainty due to the lepton energy scale is 12 (20) MeV for muon (electron) channel. We expect to get the ultimate combined Tevatron measurement with an uncertainty of 15 MeV.

W boson production charge asymmetry. At the Tevatron, W^+ (W^-) bosons are produced primarily by the annihilation of u (d) quarks in the proton with d ($\bar{u}$) quarks in the antiproton. Due to the fact that u quarks in the proton on average carry more momentum than d quarks, W^+ (W^-) boson tends to move along the proton (antiproton) direction, and thus results in an asymmetry in the W boson rapidity distribution between W^+ and W^- boson production. This asymmetry is sensitive to the PDFs which describes the fraction of momentum carried by each parton (x) in the proton.

Since we can not measure the longitudinal momentum of the neutrino in $W \rightarrow \ell\nu$ decays, the W charge asymmetry is traditionally measured as a function of the decay lepton pseudorapidity (η). The DØ collaboration analyzed 0.75 fb^{-1} of data in electron channel and measured electron charge asymmetry for events with $E_T^e > 25 \text{ GeV}$, $p_T^e > 25 \text{ GeV}$ and electron $|\eta| < 3.2$ [3]. The asymmetry is also measured in two electron E_T bins to probe partons in different x regions. The detector effects corrected charge asymmetry is compared to a NLO perturbative QCD calculation with CTEQ6.6 and MRST04 PDF sets. The measured charge asymmetries tend to be lower than the theoretical predictions for high rapidity electrons.

In a recent analysis using 1 fb^{-1} of data by the CDF collaboration [4], the W boson charge asymmetry is measured directly for the first time. The longitudinal momentum of the neutrino is estimated on an event-by-event basis using the W boson mass constraint. The measured asymmetry is found to have good agreement with the predictions of a

NNLO QCD calculation using the MRST 2006 NNLO PDF sets and a NLO QCD calculation using the CTEQ6.1 NLO PDF sets.

For both measurements, except the largest rapidity bin ($2.8 < |\eta| < 3.2$) in the $D\bar{O}$ measurement, the overall experimental uncertainties are smaller than the uncertainties due to PDFs. These measurements can be used to make tighter constraints on the global PDFs fits and reduce the uncertainty on the future W boson mass measurements.

DIBOSON PRODUCTION

The diboson production at the Tevatron is sensitive to the couplings between gauge bosons. These triple gauge couplings (TGCs) are a direct consequence of the non-Abelian group structure of the SM. Understanding diboson production is also critical for Higgs searches where dibosons are a major source of backgrounds in several important channels. The diboson measurements at the Tevatron often involve measurements of the overall production cross section and TGCs. The diboson processes with one charged W boson in the final state are discussed here. The general Lorentz invariant effective Lagrangian describing WWV ($V = \gamma$ or Z) vertices has five independent parameters with the assumption of EM gauge invariance and C and P conservation. The five parameters are g_1^Z , κ_Z , κ_γ , λ_Z and λ_γ . In the SM, $g_1^Z = \kappa_{Z,\gamma} = 1$ and $\lambda_{Z,\gamma} = 0$. These couplings are often written in terms of their deviations from their SM values as $\Delta g_1^Z = g_1^Z - 1$ and $\Delta \kappa_{Z(\gamma)} = \kappa_{Z(\gamma)} - 1$. A dipole form factor $\alpha(\hat{s}) = \alpha_0/(1 + \hat{s}/\Lambda^2)^2$ for one arbitrary coupling α is introduced to avoid the unitarity violation, where Λ is the new physics energy scale, and the limits are set in terms of α_0 .

$WW \rightarrow \ell\nu\ell\nu$. Both CDF and $D\bar{O}$ Collaborations select WW events with two high p_T leptons and large $\cancel{E}_T$. Using 1 fb^{-1} of data [5], $D\bar{O}$ observed $22(ee)$, $64(e\mu)$ and $14(\mu\mu)$ candidates. These numbers are consistent with the SM predictions of $23.5 \pm 1.9(ee)$, $68.6 \pm 3.9(e\mu)$ and $10.8 \pm 0.6(\mu\mu)$ events. The measured cross section is $11.5 \pm 2.1(\text{stat+syst}) \pm 0.7(\text{lumi}) \text{ pb}$, which is consistent with the SM prediction of $12.4 \pm 0.8 \text{ pb}$. A similar measurement by CDF used 3.6 fb^{-1} of data [6]. This measurement makes use of matrix element (ME) based likelihood ratios. Event kinematics are used to assign an event-by-event probability $\mathbf{P}$ based on leading-order ME cross section calculation for the WW process and for the major background processes such as WW , $W\gamma$ and W +jet. The background fractions are estimated using the GEANT MC simulation, while the signal fraction is extracted using a likelihood fit to the LR_{WW} distribution. LR_{WW} is defined as $P_{WW}/(P_{WW} + \sum_i k_i P_i)$ with k_i as the relative fraction of each background source, P_{WW} and P_i represent the event probability for WW and each background process respectively. The measured cross section is $12.1 \pm 0.9(\text{stat})_{-1.4}^{+1.6}(\text{syst}) \text{ pb}$, which is again consistent with the SM prediction. $D\bar{O}$ also used the leading and trailing lepton p_T s to set limits on aTGCs assuming $\Lambda = 2 \text{ TeV}$. The results are $-0.14 < \Delta g_1^Z < 0.30$, $-0.54 < \Delta \kappa_\gamma < 0.83$ and $0.14 < \lambda_Z = \lambda_\gamma < 0.19$ under the $SU(2)_L \times U(1)_Y$ -conserving constraints, and $-0.12 < \Delta \kappa_Z = \Delta \kappa_\gamma < 0.35$, with the same $\lambda_{Z(\gamma)}$ limits as above, under the $WW\gamma = WWZ$ constraints.

$WW/WZ \rightarrow \ell\nu jj$. The DØ Collaboration recently reported the first evidence of $WW/WZ \rightarrow \ell\nu jj$ process at the hadron collider [7]. The signatures are one high p_T lepton, large $\cancel{E}_T$ and two high p_T jets. The background arises mainly due to $W/Z +$ jets, QCD multijet, and $t\bar{t}$ events. A multivariate classifier (Random Forest) is used to separate the signal from backgrounds. The signal and background contents are determined by fitting the signal and background random forest templates to the data. The measured cross section is $20.2 \pm 2.5(\text{stat}) \pm 3.6(\text{syst}) \pm 1.2(\text{lumi})$ pb, consistent with the SM prediction of 16.1 ± 0.9 pb. The dijet p_T spectrum is used to set aTGCs limits assuming $\Lambda = 2$ TeV: $-0.12 < \Delta g_1^Z < 0.19$, $-0.44 < \Delta \kappa_\gamma < 0.55$ and $-0.10 < \lambda_{Z,\gamma} < 0.11$ under the $SU(2)_L \times U(1)_Y$ -conserving constraints, and $-0.16 < \Delta \kappa_{Z,\gamma} < 0.23$ and $-0.11 < \lambda_{Z,\gamma} < 0.11$, under the $WW\gamma = WWZ$ constraints.

$WZ \rightarrow jj\ell\ell$. The CDF Collaboration also reported a search for anomalous WZ production at the Tevatron using the two charged leptons, two jets final state [8]. Three bins in dilepton p_T are used: 105 – 140 GeV (control region), 140 – 210 GeV (medium region) and > 210 GeV (high region). Events in the control region are mainly used to validate data modeling and determine backgrounds. The dijet mass M_{jj} for the medium and high regions are used to set limits on the cross section and aTGCs. The 95% CL upper limit on the cross section is found to be 234 fb and 135 fb using events in the medium and high region respectively. Under the $WW\gamma = WWZ$ constraints, the limits on the aTGCs are $-0.20 < \Delta g_1^Z < 0.29$, $-1.01 < \Delta \kappa_{Z,\gamma} < 1.27$ and $-0.16 < \lambda_{Z,\gamma} < 0.17$ assuming $\Lambda = 2$ TeV, and $-0.22 < \Delta g_1^Z < 0.32$, $-1.09 < \Delta \kappa_{Z,\gamma} < 1.40$ and $-0.18 < \lambda_{Z,\gamma} < 0.18$ assuming $\Lambda = 1.5$ TeV.

CONCLUSIONS

With the increasingly large datasets, both CDF and DØ continue to improve our understanding of electroweak production at the hadron collider.

ACKNOWLEDGMENTS

I would like to thank all my CDF and DØ colleagues for their hard works, as well as the conference organizers for hosting an excellent conference.

REFERENCES

1. V.M. Abazov *et al.* (DØ Collaboration), <http://www-d0.fnal.gov/Run2Physics/WWW/results/prelim/EW/E27>.
2. T. Aaltonen *et al.* (CDF Collaboration), <http://www-cdf.fnal.gov/physics/ewk/2008/wmass/>.
3. V.M. Abazov *et al.* (DØ Collaboration), Phys. Rev. Lett. **101**, 211801 (2008).
4. T. Aaltonen *et al.* (CDF Collaboration), Phys. Rev. Lett. **102**, 181801 (2009); A. Bodek *et al.*, Phys. Rev. D **77**, 111301(R) (2008).
5. V.M. Abazov *et al.* (DØ Collaboration), submitted to Phys. Rev. Lett., arXiv:0904.0673.
6. T. Aaltonen *et al.* (CDF Collaboration), <http://www-cdf.fnal.gov/physics/ewk/wwllll/index.html>.
7. V.M. Abazov *et al.* (DØ Collaboration), Phys. Rev. Lett. **102**, 161801 (2009).
8. T. Aaltonen *et al.* (CDF Collaboration), <http://www-cdf.fnal.gov/physics/ewk/2008/WWZaTGC/>.